

Fake CPT Violation in Disappearance Neutrino Oscillations

Zhi-zhong Xing

*Institute of High Energy Physics, P.O. Box 918 (4), Beijing 100039, China
(Electronic address: xingzz@mail.ihep.ac.cn)*

Abstract

We make an analysis of the fake CPT -violating asymmetries between the survival probabilities of neutrinos and antineutrinos, induced by the terrestrial matter effects, in three different scenarios of long-baseline neutrino oscillation experiments with $L = 730$ km, $L = 2100$ km and $L = 3200$ km. In particular, the dependence of those asymmetries on the Dirac-type CP -violating phase of the lepton flavor mixing matrix is examined.

PACS number(s): 14.60.Pq, 13.10.+q, 25.30.Pt

It is a real challenge in particle physics to understand the origin of neutrino masses and the mechanism of lepton flavor mixing [1]. Although the present non-accelerator neutrino oscillation experiments have yielded some impressive constraints on the lepton flavor mixing parameters [2–4], more precise measurements of those parameters are desirable and rely on a new generation of accelerator neutrino oscillation experiments with long baselines [5], including the possible neutrino factories [6]. The terrestrial matter effects in such long-baseline neutrino experiments must be taken into account, since they unavoidably modify the genuine behaviors of neutrino oscillations in vacuum. A particular manifestation of the terrestrial matter effects is that they may give rise to *fake CPT*-violating asymmetries between the survival probabilities of neutrinos and antineutrinos [7].

The purpose of this Brief Report is to illustrate the fake *CPT*-violating asymmetries between $\nu_\alpha \rightarrow \nu_\alpha$ and $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha$ transitions (for $\alpha = e, \mu, \tau$) in three scenarios of long-baseline neutrino oscillation experiments: $L = 730$ km, $L = 2100$ km and $L = 3200$ km. In particular, the dependence of those asymmetries on the Dirac-type *CP*-violating phase of the lepton flavor mixing matrix will be examined.

Let us consider a transition $\nu_\alpha \rightarrow \nu_\alpha$ (for $\alpha = e, \mu$ or τ) and its *CPT*-conjugate process $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha$ in vacuum. *CPT* invariance implies that the probabilities of $\nu_\alpha \rightarrow \nu_\alpha$ and $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha$ transitions amount to each other. Explicitly, we have

$$P(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4 \sum_{i < j} \left[|V_{\alpha i}|^2 |V_{\alpha j}|^2 \sin^2 \left(1.27 \frac{\Delta m_{ji}^2}{E} L \right) \right], \quad (1)$$

where $V_{\alpha i}$ (for $\alpha = e, \mu, \tau$ and $i = 1, 2, 3$) denote the elements of the lepton flavor mixing matrix V , $\Delta m_{ji}^2 \equiv m_j^2 - m_i^2$ with m_i being the neutrino masses (in unit of eV), L is the distance between the production and interaction points of ν_α (in unit of km), and E is the neutrino beam energy (in unit of GeV). As *T* conservation is automatic for the survival probabilities of ν_α and $\bar{\nu}_\alpha$ neutrinos, *CP* symmetry holds between $P(\nu_\alpha \rightarrow \nu_\alpha)$ and $P(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha)$ as a direct consequence of *CPT* invariance.

In a realistic long-baseline neutrino experiment, the genuine pattern of $\nu_\alpha \rightarrow \nu_\alpha$ and $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha$ transitions is modified due to the presence of terrestrial matter effects. Defining the *effective* neutrino masses $\tilde{m}_i$ and the *effective* lepton flavor mixing matrix $\tilde{V}$ in matter, one may write out the *effective* probability of $\nu_\alpha \rightarrow \nu_\alpha$ in the same form as Eq. (1):

$$\tilde{P}(\nu_\alpha \rightarrow \nu_\alpha) = 1 - 4 \sum_{i < j} \left[|\tilde{V}_{\alpha i}|^2 |\tilde{V}_{\alpha j}|^2 \sin^2 \left(1.27 \frac{\Delta \tilde{m}_{ji}^2}{E} L \right) \right], \quad (2)$$

where $\Delta \tilde{m}_{ji}^2 \equiv \tilde{m}_j^2 - \tilde{m}_i^2$. The analytically exact expressions of $\tilde{m}_i^2$ and $\tilde{V}_{\alpha i}$, which depend upon Δm_{ji}^2 , $V_{\alpha i}$ and the matter parameter $A = 2\sqrt{2} G_F N_e E$ with N_e being the background density of electrons and E being the neutrino beam energy, have been given in Ref. [8]. Note that the *effective* probability of $\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha$ can simply be read off from Eq. (2) with the replacements $V_{\alpha i} \Rightarrow V_{\alpha i}^*$ and $A \Rightarrow -A$. There must be an asymmetry between $\tilde{P}(\nu_\alpha \rightarrow \nu_\alpha)$ and $\tilde{P}(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha)$, since they are associated respectively with $(V, +A)$ and $(V^*, -A)$. This *fake CPT* asymmetry, denoted as

$$\Delta_\alpha \equiv \tilde{P}(\nu_\alpha \rightarrow \nu_\alpha) - \tilde{P}(\bar{\nu}_\alpha \rightarrow \bar{\nu}_\alpha), \quad (3)$$

arises purely from the terrestrial matter effects. A measurement of Δ_α is expected to shed light on both the matter parameter A and the neutrino mixing parameters Δm_{ji}^2 and $V_{\alpha i}$.

To carry out numerical calculations of the asymmetry Δ_α , we adopt the standard parametrization for the lepton flavor mixing matrix V [9]:

$$V = \begin{pmatrix} c_1 c_3 & s_1 c_3 & s_3 \\ -c_1 s_2 s_3 - s_1 c_2 e^{-i\delta} & -s_1 s_2 s_3 + c_1 c_2 e^{-i\delta} & s_2 c_3 \\ -c_1 c_2 s_3 + s_1 s_2 e^{-i\delta} & -s_1 c_2 s_3 - c_1 s_2 e^{-i\delta} & c_2 c_3 \end{pmatrix} \begin{pmatrix} 1 & 0 & 0 \\ 0 & e^{i\rho} & 0 \\ 0 & 0 & e^{i\sigma} \end{pmatrix} \quad (4)$$

with $s_i \equiv \sin \theta_i$ and $c_i \equiv \cos \theta_i$ (for $i = 1, 2, 3$). Since normal neutrino oscillations are completely insensitive to the Majorana-type CP -violating phases ρ and σ , we only need inputs of the other four parameters $\theta_1, \theta_2, \theta_3$ and δ . Indeed $(\theta_1, \theta_2, \theta_3)$ are related respectively to the mixing angles of solar, atmospheric and reactor disappearance neutrino oscillations, at least in the leading-order approximation. As the large-angle Mikheyev-Smirnov-Wolfenstein (MSW) solution to the solar neutrino problem [10] is most favored by the latest Super-Kamiokande and SNO data [11], we typically take $\theta_1 = 35^\circ$ together with $\Delta m_{21}^2 \approx \Delta m_{\text{sun}}^2 \approx 5 \cdot 10^{-5} \text{ eV}^2$. We take $\theta_2 = 40^\circ$ and $\Delta m_{31}^2 \approx \Delta m_{\text{atm}}^2 \approx 3 \cdot 10^{-3} \text{ eV}^2$ in view of the present data on atmospheric neutrino oscillations [2], as well as $\theta_3 = 6^\circ$ in view of the upper bound $\sin^2 2\theta_3 < 0.1$ set by the CHOOZ and Palo Verde Collaborations [3]. The size of the Dirac-type CP -violating phase δ remains experimentally unrestricted. We consider two distinct possibilities for illustration: $\delta = 0^\circ$ (with vanishing CP violation) and $\delta = 90^\circ$ (with maximal CP violation).

If the baseline of neutrino oscillations is about 3000 km or shorter, it is a good approximation to assume the constant matter density of the earth's crust [12]: $A \approx 2.2 \cdot 10^{-4} \text{ eV}^2 E / [\text{GeV}]$ with E being the neutrino beam energy. Here let us consider three interesting scenarios of long-baseline neutrino experiments: $L = 730 \text{ km}$, $L = 2100 \text{ km}$ and $L = 3200 \text{ km}$. The first baseline corresponds essentially to a neutrino source at the Fermilab pointing toward the Soudan mine or that at CERN toward the Gran Sasso underground laboratory; the second baseline is compatible with a possible high-intensity neutrino beam from the High Energy Proton Accelerator in Tokaimura to a detector located in Beijing; and the third baseline could be a favorable choice for the future neutrino factories to measure leptonic CP - and T -violating asymmetries [13]. We are then able to calculate the fake CPT -violating asymmetries Δ_e , Δ_μ and Δ_τ with the help of the formulas presented already in Ref. [8] and the inputs given above. Our numerical results are shown in Figs. 1, 2 and 3. Two comments are in order.

(a) Different from Δ_μ and Δ_τ , Δ_e is independent of the value of δ . This feature can easily be understood as follows. $\tilde{P}(\nu_e \rightarrow \nu_e)$ relies on $\tilde{V}_{e1}$, $\tilde{V}_{e2}$ and $\tilde{V}_{e3}$, which are in turn dependent on V_{e1} , V_{e2} and V_{e3} but independent of the other matrix elements of V [8]. Since V_{e1} , V_{e2} and V_{e3} have nothing to do with δ in the afore-mentioned standard parametrization, $\tilde{P}(\nu_e \rightarrow \nu_e)$ is completely independent of δ . A similar argument holds for $\tilde{P}(\bar{\nu}_e \rightarrow \bar{\nu}_e)$. Therefore the asymmetry Δ_e keeps unchanged for arbitrary values of δ .

(b) For the chosen range of E (i.e., $1 \text{ GeV} \leq E \leq 20 \text{ GeV}$), the magnitude of Δ_e at the maximal resonance can be as large as 2.7% for $L = 730 \text{ km}$, 7.5% for $L = 2100 \text{ km}$, or 11% for $L = 3200 \text{ km}$. A measurement of Δ_e is therefore possible, at least in principle. The sizes of Δ_μ and Δ_τ are too small to be detectable for $L = 730 \text{ km}$; but they can be as large as a

few percent at the resonances, if $L \geq 2000$ km and $\delta \sim 0^\circ$. Therefore the dependence of Δ_μ and Δ_τ on δ might be detectable.

We find that the behavior of Δ_α is quite stable, even if the matter parameter A fluctuates slightly around the originally chosen value. The results obtained above are essentially unchanged if Δm_{21}^2 flips its sign, but they are sensitive to the sign flip of Δm_{31}^2 . In particular, the relation $\Delta_e(-\Delta m_{31}^2) \approx -\Delta_e(+\Delta m_{31}^2)$ is found to hold approximately. Therefore the measurement of Δ_α may help determine the relative sizes of m_2 and m_3 , which are important to fix the neutrino mass spectrum. We have also noticed that Δ_α is sensitive to θ_3 [14]. If θ_3 were too small (e.g., $\theta_3 < 1^\circ$), the asymmetries Δ_e , Δ_μ and Δ_τ would be too small to be practically detected. In a variety of phenomenological models for lepton masses and flavor mixing [1,15], $\theta_3 \sim \arctan \sqrt{m_e/m_\mu} \approx 4^\circ$ is expected to hold. The measurement of Δ_α may provide a valuable chance to determine or constrain the smallest flavor mixing angle θ_3 in the lepton sector.

Finally it is worth mentioning that there might be small *genuine* CPT -odd interactions in neutrino oscillations [16], and they might be entangled with the fake CPT -odd interactions induced by the terrestrial matter effects. Whether the former can be disentangled from the latter in realistic long-baseline neutrino oscillation experiments remains an open question and deserves a systematic study.

The author is grateful to L. Wolfenstein and L.Y. Shan for useful discussions.

REFERENCES

- [1] For recent reviews with extensive references, see: G. Altarelli and F. Feruglio, Phys. Rep. **320**, 295 (1999); R.N. Mohapatra, hep-ph/9910365 (to appear in *Current Aspects of Neutrino Physics*, edited by D. Caldwell, Springer-Verlag); H. Fritzsch and Z.Z. Xing, Prog. Part. Nucl. Phys. **45**, 1 (2000); S.M. Barr and I. Dorsner, Nucl. Phys. B **585**, 79 (2000); P. Langacker, Nucl. Phys. Proc. Suppl. **100**, 383 (2001).
- [2] Y. Fukuda *et al.*, Phys. Rev. Lett. **81**, 1562 (1998); *ibid.* **81**, 4279 (1998); <http://www-sk.icrr.u-tokyo.ac.jp/dpc/sk/>.
- [3] CHOOZ Collaboration, M. Apollonio *et al.*, Phys. Lett. B **420**, 397 (1998); Palo Verde Collaboration, F. Boehm *et al.*, Phys. Rev. Lett. **84**, 3764 (2000).
- [4] SNO Collaboration, Q.R. Ahmad *et al.*, nucl-ex/0106015.
- [5] P. Fisher, B. Kayser, and K.S. McFarland, Ann. Rev. Nucl. Part. Sci. **49** (1999) 481; A. de Santo, hep-ph/0106089; and references therein.
- [6] B. Autin *et al.*, CERN 99-02 (1999); D. Ayres *et al.*, physics/9911009; C. Albright *et al.*, hep-ex/0008064; and references therein.
- [7] N. Cabibbo, Phys. Lett. B **72**, 333 (1978); V. Barger, K. Whisnant, and R.J.N. Phillips, Phys. Rev. Lett. **45**, 2084 (1980); J. Bernabéu, hep-ph/9904474.
- [8] Z.Z. Xing, Phys. Lett. B **487**, 327 (2000); Phys. Rev. D **64**, 073014 (2001).
- [9] H. Fritzsch and Z.Z. Xing, Phys. Lett. B **517**, 363 (2001).
- [10] L. Wolfenstein, Phys. Rev. D **17**, 2369 (1978); S.P. Mikheyev and A. Yu Smirnov, Yad. Fiz. (Sov. J. Nucl. Phys.) **42**, 1441 (1985).
- [11] J. Goodman (Super-Kamiokande Collaboration), talk given at the 20th International Symposium on Lepton and Photon Interactions at High Energies, July 2001, Rome, Italy; J. Klein (SNO Collaboration), talk given at the same conference.
- [12] V. Barger, S. Geer, R. Raja, and K. Whisnant, Phys. Rev. D **62**, 013004 (2000); I. Mocioiu and R. Shrock, Phys. Rev. D **62**, 053017 (2000); T. Ota and J. Sato, Phys. Rev. D **63**, 093004 (2001).
- [13] See, e.g., A. Romanino, Nucl. Phys. B **574**, 675 (2000); A. Cervera *et al.*, Nucl. Phys. B **579**, 17 (2000); V. Barger, S. Geer, R. Raja, and K. Whisnant, Phys. Rev. D **62**, 073002 (2000); Z.Z. Xing, Phys. Rev. D **63**, 073012 (2001); S.J. Parke and T.J. Weiler, Phys. Lett. B **501**, 106 (2001); I. Mocioiu and R. Shrock, hep-ph/0106139; L.Y. Shan, B.L. Young, and X. Zhang, hep-ph/0110414; Y.F. Wang, K. Whisnant, Z. Xiong, J.M. Yang, and B.L. Young, hep-ph/0111317.
- [14] M.C. Bañuls, G. Barenboim, and J. Bernabéu, hep-ph/0102184.
- [15] H. Fritzsch and Z.Z. Xing, Phys. Lett. B **372**, 265 (1996); Phys. Lett. B **440**, 313 (1998); Phys. Rev. D **61**, 073016 (2000); Z.Z. Xing, Phys. Rev. D **64**, 093013 (2001).
- [16] S. Coleman and S.L. Glashow, Phys. Rev. D **59**, 116008 (1999); V. Barger, S. Pakvasa, T.J. Weiler, and K. Whisnant, Phys. Rev. Lett. **85**, 5055 (2000).

FIGURES




FIG. 1. Illustrative plots for fake CPT -violating asymmetries Δ_e , Δ_μ and Δ_τ .




FIG. 2. Illustrative plots for fake CPT -violating asymmetries Δ_e , Δ_μ and Δ_τ .




FIG. 3. Illustrative plots for fake CPT -violating asymmetries Δ_e , Δ_μ and Δ_τ .